



BROAD SPECTRUM FUNGICIDE/BACTERICIDE For foliar control of bacterial and fungal plant diseases

SPECIMEN LABEL

ACTIVE INGREDIENTS:

Sodium Carbonate Peroxyhydrate85.00%

OTHER INGREDIENTS: 15.00%

TOTAL: 100.00%

*Contains 27.60% Hydrogen Peroxide by weight.

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

First Aid

If in eyes

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes.
- Call a poison control center or doctor for treatment advice.

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center.
- Do not give anything by mouth to an unconscious person.

If inhaled

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: CORROSIVE. Causes irreversible eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Handlers who may be exposed to the dilute through application or other tasks must wear: long-sleeved shirt and long pants, and shoes plus socks. Handlers who may be exposed to the concentrate through mixing, loading, application, or other tasks must wear: coveralls over long-sleeved shirt and long pants, rubber gloves, chemical resistant footwear plus socks, and protective eyewear (goggles or face shield). Follow manufacturer's instructions for cleaning / maintaining PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. If

Net Contents:

50 lbs.

EPA Registration No. 70299-15 • V15

Sold by BioSafe Systems LLC

22 Meadow Street, East Hartford, CT 06108

1-888-273-3088 (toll-free) www.biosafesystems.com

V15



no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the out side of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. This product is highly toxic to bees and other beneficial insects exposed to direct contact on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Do not apply this product or allow it to drift to crops where beneficials are part of an integrated pest management strategy. Do not use near waterbody margins during amphibian breeding seasons.

PHYSICAL AND CHEMICAL HAZARDS

Strong oxidizing agent. **Corrosive.** Do not bring in contact with other pesticides, cleaners or oxidative agents.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the state or tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), notification to workers, and Restricted-Entry Interval (REI). The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

For enclosed environments

There is a restricted entry of one (1) hour for this product when applied via fogging or spraying to growing plants, surfaces, equipment, structures and non-porous surfaces in enclosed environments such as glasshouses and greenhouses. PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls worn over long-sleeved shirt and pants, waterproof gloves and shoes plus socks.

There is a restricted entry of zero (0) hours for pre-plant dip, seed treatment, soil drench, mop, sponge, dip, soak, rinse or other non-spraying or non-fogging application methods when used in enclosed environments such as glasshouses and greenhouses.

For field applications

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

This product is not intended as treatment against any public health organism for any use on this label.

Do not allow spray to drift from the application site and contact people or structures that people occupy.

FOLIAR SPRAY TREATMENTS

Use PerCarb as a contact foliar fungicide/bactericide to treat and control plant pathogenic diseases on field grown crops, tree crops, berries, small fruits, vine crops, or crops grown in similar sites.

PerCarb may be applied up to and including the day of harvest.

COMPATIBILITY

PerCarb is compatible as a tank-mix with many commonly used pesticides, fertilizers, adjuvants and non-ionic surfactants but has not been fully evaluated with all of these. Do not tank mix PerCarb in spray tanks with pesticides, surfactants or fertilizers before conducting a compatibility test to show it is physically compatible, effective and non-injurious under your use conditions.

To ensure compatibility, evaluate them prior to use as follows: Using a suitable container, add proportional amounts of product to water. Add wettable powders first, followed by water dispersible granules, then by liquid flowables and lastly, emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible. Excessive bubbling and/or pressure are indications of incompatibility. Test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

PLANT SENSITIVITY TESTING

Be sure to use PerCarb at labeled dilutions as solutions more concentrated can result in leaf necrosis for some crops (i.e., do not use at concentrations greater than labeled rates). PerCarb provides a balanced source of the active ingredient directly to the plant surface. PerCarb has been used and tested on many varieties of plant material; however, the nature of the target plant, environmental conditions, plant vigor and the use of other pesticides can all affect plant sensitivity to PerCarb. Therefore, before treating large numbers of plants, test PerCarb on a few plants for sensitivity.

Application of PerCarb for curative control of obligate organisms living in the plant tissue (such as Downy and Powdery Mildew) can result in lesions on plant tissue. PerCarb will oxidize parasitic organisms living in plant tissue that are not always visible to the naked eye. Resulting oxidative effects can include spotting, or drying of the plant tissue where organisms inhabited tissue.

MIXING INSTRUCTIONS

PerCarb will go into solution with clean water at no more than 1/2 pound per gallon of water. Dissolution of PerCarb in cold water takes approximately 5 minutes, less in warm water. When mixing, fill the spray tank one-half full of water,

add PerCarb slowly to the clean tank, while hydraulic and or mechanical agitation is operating. The additions of any spreaders should be added only after they have been predetermined to be chemically compatible with PerCarb. PerCarb works best when diluted with water containing low levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing.

Before tank mixing with fertilizers, fungicides, or bactericides, conduct a compatibility test for each combination. Make a test solution and shake or stir vigorously. Excessive bubbling and/or pressure are indications of incompatibility.

Do not mix PerCarb with other products with an acidic pH. Do not adjust pH after mixing of product. Acidification of solution will cause reduced product performance.

PerCarb is a strong oxidizing agent and may react with residues of metal-based fungicides or supplements. Do not apply PerCarb as a foliar spray immediately following foliar applications of metal-based products. Allow at least 24 hrs. after application of metal-based products before applying PerCarb as a foliar spray. Check the label of the metal-based product prior to application for specific instructions for use with other fungicide products.

Note: Use spray solutions within two hours of preparation, do not store and reuse mixed spray solution.

FOLIAR SPRAY APPLICATION INSTRUCTIONS

Apply as a foliar spray throughout the growing season on the listed crops and diseases.

Apply using backpack, hand-held, high volume, and boom or ground sprayer. For best results, use a properly calibrated sprayer or similar equipment. Check equipment, and calibrate frequently.

PerCarb works by surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wet. PerCarb does not produce a distinct odor or deleterious effects to plants when used in accordance with label directions. Do not use at stronger than recommended rates, as leaf burn may result.

Prepare a solution by mixing 0.25–0.75 lbs. (4–12 oz.) of PerCarb per 25 gallons of water (1–3 lbs. per 100 gallons of water). Use up to 400 gallons of spray solution per acre. For the most effective results, use enough volume of spray solution to obtain complete and uniform coverage of foliage and stems. Spray solution volume per acre will vary depending on crop type and stage of development.

FOLIAR APPLICATION TIMING

Begin preventative applications at low rates when conditions are favorable for disease development such as rainy weather or during periods of high relative humidity. Maintain a 7–10 day spray interval. Begin using high rate at first sign of disease. Maintain a 7-day spray interval until control is achieved

EARLY AND LATE DORMANT SPRAY APPLICATION INSTRUCTIONS:

Use as a dormant spray for early and late season applications on tree nuts, tree fruits, small fruits, berries and vine crops prior to bud opening.

Prepare a solution by mixing 0.75–1.0 lbs. of PerCarb per 25 gallons of water (3–4 lbs. per 100 gallons of water). Use up to 400 gallons of spray solution per acre. For the most effective results, use enough volume of spray solution to obtain complete and uniform coverage.

PerCarb Dilution Rate lbs./25 gallons of water	Lbs. of PerCarb per Acre			
	Total Spray Volume (Gallons/Acre)			
	50	100	200	400
0.25	0.50	1	2	4
0.50	1.0	2	4	8
0.75	1.5	3	6	12
1.0 (Dormant Sprays Only)	2	4	8	16

Note: For sensitive crops, such as spinach and swiss chard, do not exceed concentrations greater than 1 lb. /100 gallon of water and increase spray frequency.

- Do not apply this product through any type of irrigation system.
- Do not apply directly to soil as a directed spray or drench.

CROPS AND DISEASES APPLICATION CHART (ALPHABETICAL BY CROP GROUPING)

Crop	Diseases	Lbs./100 gallons of water
Asparagus	Asparagus Rust	Foliar: 1–3 lbs.
Berries: Blackberry, Blueberry Cranberry, Raspberry	Alternaria, Botrytis, Downy Mildew, Mummy Berry Leaf Blight, Powdery Mildew, Fruit Rot, Rust	Foliar: 1–3 lbs.
		Dormant 3–4 lbs. Apply in early and late dormancy prior to bud break. Do not apply to blooming crops.
Bulb Vegetables: Garlic, Green Onions, Leeks, Onions, Scallions, Shallots	Alternaria Leaf Blight, Bacterial Leaf Blight, Botrytis Rot, Downy Mildew, Powdery Mildew	Foliar: 1–3 lbs.
Cereal Grains & Commodities: Barley, Corn (field), Millet, Oats Popcorn, Rice, Rye, Sorghum (Milo) Soybeans, Sweet Corn, Wheat, Wild Rice	Anthracnose, Bacterial Leaf Blight, Blast, Brown Leaf Spot, Common Rust, Common Smut, Downy Mildew, Leaf Smut, Sheath Blight	Foliar: 1–3 lbs.
Citrus Crops: Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Orange, Tangerine	Alternaria, Anthracnose, Brown Rot, Citrus Canker, Powdery Mildew, Rust, Scab	Foliar: 1–3 lbs.
		Dormant: 3–4 lbs. Apply in early and late dormancy prior to bud break. Do not apply to blooming crops.
Coffee	Bacterial Blight, Leaf Rust	Foliar: 1–3 lbs.
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale	Black Rot, Downy Mildew, Powdery Mildew	Foliar: 1–3 lbs.
Cotton	Angular Leaf Spot	Foliar: 1–3 lbs.
Cucurbit Crops: Cucumber, Melons, Pumpkin, Squash	Alternaria leaf spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Leaf Spot, Powdery Mildew	Foliar: 1–3 lbs.
Forage and Fodder Crops: Alfalfa Hay, Clover, Trefoil, Lupine, Grasses	Cerospora Leaf Spot	Foliar: 1–3 lbs.
Fruiting Vegetables: Eggplant, Peppers, Tomatoes, Tomatillos	Anthracnose, Bacterial Spot, Bacterial Speck , Early Blight, Late Blight	Foliar: 1–3 lbs.
Globe Artichokes	Black Rot, Botrytis Blight/Grey Mold, Crown Rot, Powdery Mildew	Foliar: 1–3 lbs.
Grapes	Botrytis, Downy Mildew, Powdery Midew	Foliar: 1–3 lbs.
		Dormant: 3–4 lbs. Apply in early and late dormancy prior to bud break. Do not apply to blooming crops.
Herbs and Spices: Basil, Chives, Cilantro, Coriander, Dill, Mint, Rosemary, Sage	Anthracnose, Downy Mildew, Powdery Mildew	Foliar: 1–3 lbs.
		Special Note: For sensitive crops do not exceed concentrations greater than 1 lb./100 gallons of water and increase spray frequency.
Hops	Downy Mildew, Powdery Mildew	Foliar: 1–3 lbs.
Leafy Vegetables: Arugula, Celery, Chickory Root, Endive, Fennel, Lettuce, Spinach, Rhubarb, Radicchio, Swiss Chard	Downy Mildew, Early Blight, Powdery Mildew	Foliar: 1–2 lbs.
		Special Note: For sensitive crops, do not exceed concentrations greater than 1 lb./100 gallons of water and increase spray frequency.
Legumes: Chick Peas, Dry Beans, Lima Beans, Peas, Snap Beans, Soybeans	Anthracnose, Bacterial Blight, Powdery Mildew, Rust, White Mold	Foliar: 1–3 lbs.
Peanuts	Rust, Leaf Spot	Foliar: 1–3 lbs.
Pome Fruits: Apples, Pears, Loquats, Mayhaws, Quince	Flyspeck, Powdery Mildew, Rusts, Scab, Sooty Mold	Foliar: 1–3 lbs.
		Dormant: 3-4 lbs. Apply in early and late dormancy prior to bud break. Do not apply to blooming crops
Root & Tuber Vegetables: Artichokes, Beets, Carrots, Ginseng, Horseradish, Parsnip, Potatoes, Radish, Rutabaga, Sugar Beets, Sweet Potatoes, Taro, Turnips, Yams	Alternaria Leaf Spot/Blight, Bacterial Leaf Spot/ Blight, Early Blight, Late Blight, Leaf Spot Powdery Mildew, Potato Brown Rot, White Mold	Foliar: 1–3 lbs.
Stone Fruits: Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	Botrytis Blight, Brown Rot / Blossom Blight, Cherry Leaf Spot, Leaf Spot, Powdery Mildew Rust, Scab	Foliar: 1–3 lbs.
		Dormant: 3–4 lbs. Apply in early and late dormancy prior to bud break. Do not apply to blooming crops.

Strawberries	Anthrachnose, Angular Leaf Spot, Botrytis/Fruit Rot, Powdery Mildew	Foliar: 1–3 lbs.
Tobacco (Field/Float Beds)	Blue Mold	Foliar: 1–3 lbs.
Tree Nuts: Almonds, Brazil Nuts, Cashews, Filberts, Macadamias, Pecans, Pistachios, Walnuts	Alternaria Leaf Spot/Blight, Anthracnose, Brown Rot, Bacterial Blight, Bacterial Canker, Jacket Rot, Walnut Blight, Leaf Spot, Leaf Blight, Filbert Blight, Powdery Mildew, Botryosphaeria Blight, Phomopsis Blight, Rust, Scab	Foliar: 1–3 lbs. Dormant: 3–4 lbs. Apply in early and late dormancy prior to bud break. Do not apply to blooming crops
Tropical Fruit: Avocado, Casaba, Coconut, Dates, Guava, Kiwi, Mango, Papaya, Passion Fruit, Poi, Star Fruit	Alternaria, Anthracnose, Bacterial Leaf Spot, Black Spot, Leaf Blight, Powdery Mildew, Sooty Mold	Foliar: 1–3 lbs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a cool, well-vented area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

PESTICIDE HANDLING: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited. If wastes cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Plastic Containers, All sizes: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and reclose. Shake or stir for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke

50 lb. Bags/Sacks: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty sack [bag] in application equipment. Then dispose of empty sack [bag] in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BIOSAFE SYSTEMS MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, in no event shall BIOSAFE SYSTEMS or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF BIOSAFE SYSTEMS AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF BIOSAFE SYSTEMS OR SELLER, THE REPLACEMENT OF THE PRODUCT.

BIOSAFE SYSTEMS and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of BIOSAFE SYSTEMS.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. To the extent consistent with applicable law, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of BIOSAFE SYSTEMS LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold BIOSAFE SYSTEMS and Seller harmless for any claims relating to such factors. BIOSAFE SYSTEMS warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or BIOSAFE SYSTEMS, and Buyer and User assume



For additional information on PerCarb®, call us toll-free at 1.888.273.3088 or visit www.biosafesystems.com.

©2017 BioSafe Systems, LLC. PerCarb® is a registered trademark of BioSafe Systems, LLC. Always read and follow label directions. V15-111516