



OxiDate® 5.0

FIFRA SECTION 2(e) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(e) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

ISSUE DATE:

June 16, 2022

EXPIRATION DATE:

December 31, 2025

**FOR USE AND
DISTRIBUTION ONLY
IN THE STATE OF:**

IOWA

MINNESOTA

NORTH DAKOTA

SOUTH DAKOTA

WISCONSIN

EPA REGISTRATION NO. 70299-28

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Crop: Legume Vegetables (succulent or dried) Including, but not limited to: Dry Beans

Disease: White Mold

Application Rates and Directions: OxiDate 5.0 works immediately on contact for control of plant diseases – see **Application Rates and Directions Chart**. Good coverage and wetting of the foliage is required. Apply OxiDate 5.0 in solution quantities (gallons per acre) great enough to ensure uniform coverage of upper and lower foliage, stems, branches and stalks. Final spray solution volume will depend on crop type, canopy size and/or growth stage. For increased coverage and penetration of spray, use a compatible non-ionic wetting agent/surfactant. For drift reduction and to aid spray deposition, the use of a proper adjuvant is recommended. Do not spray OxiDate 5.0 during conditions of intense plant stress. Before widespread use, run a plant sensitivity test when considering use of spray concentrations greater than 0.39% v/v (1:256) by following instructions under **Plant Sensitivity Testing**.

See **Application Rates and Directions Chart** for disease list and additional instructions.

Begin applications of OxiDate 5.0 when conditions favor the development of disease and/or during growth stages most susceptible to infection, then continue use in 3-21 day intervals or as needed. Under moderate to severe disease pressure/conditions or at first sign/symptom of disease, reduce intervals and increase rates of application until control is achieved. OxiDate 5.0 can be applied at dilution rates of 1:500-1:100 (26-128 fl. oz. per 100 gallons of water) depending on the crop group. See **Applications Rates and Directions Chart** for additional crop specific rates and instructions. Diluting OxiDate 5.0 in clean water by 75% before adding tank-mix partners and/or using a lower dilution rate, such as 1:500 (26 fl. oz. per 100 gallons of water), may improve compatibility with potential reactive tank-mix partners. See **Solution Preparation, Compatibility, and Plant Sensitivity Testing** for additional information and instructions.



BioSafe Systems, LLC

22 Meadow Street
East Hartford, CT 06108

www.BioSafeSystems.com

1-888-273-3088

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(e) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.