



OxiDate® 5.0

FIFRA SECTION 2(ee) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

ISSUE DATE:

November 10, 2024

EXPIRATION DATE:

December 31, 2027

**FOR USE AND
DISTRIBUTION ONLY
IN THE STATE OF:**

TEXAS

EPA REGISTRATION NO. 70299-28

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Crop: Strawberry

Disease: Pestalotia Leaf Spot and Fruit Rot

Causal agent: *Neopestalotiopsis* sp.

Pre-Plant Dip or Spray: Dilute at a 1:256 rate (5.0 fl. oz. per every 10 gallons of water). Thoroughly wet transplants by dipping or spraying prior to planting. Excessive foaming or bubbling during the dipping process is an indication of high levels of disease contamination. Remove dead or dying foliage prior to dipping.

Setting Water Applications: Apply at a 1:100-1:500 dilution rate (128-26 fl. oz. per 100 gallons of transplant water or starter fertilizer). Make in-furrow or dibble application at the time of plant set. OxiDate 5.0 is chemically compatible with most water-soluble fertilizers, but conduct a compatibility test prior to mixing.

Foliar applications immediately post planting: Apply at a 1:256 dilution (50 fl. oz. for 100 gallons of water). Make application immediately following planting. Apply in a sufficient amount of water to achieve runoff to soil or plastic, typically 30 to 100 gallons of spray solution per acre.

Foliar applications to established plants: Follow: "Application Rates and Directions." When conditions favor development of disease, or at first sign/symptom of disease, apply at 1:500-1:256 dilution. Maintain a 3-10 day spray schedule until control is achieved.



BioSafe Systems, LLC

22 Meadow Street
East Hartford, CT 06108

www.BioSafeSystems.com

1-888-273-3088

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.