

## FIFRA SECTION 2(ee) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

#### **ISSUE DATE:**

June 30, 2025

### **EXPIRATION DATE:**

December 31, 2028

# FOR USE AND DISTRIBUTION ONLY IN THE STATE OF:

**TEXAS** 

**EPA REGISTRATION NO.** 70299-28

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Crop: Strawberries** 

Disease(s): Phomopsis Leaf Blight and Fruit Rot\*, Pestalotia Leaf Spot and Fruit Rot\*\*

**Pre-plant dip or spray:** Dilute at a 1:100 rate (12.8 fl. oz. per every 10 gallons of water). Thoroughly wet transplants by dipping or spraying prior to planting. Excessive foaming or bubbling during the dipping process is an indication of high levels of disease contamination. Remove dead or dying foliage prior to dipping.

**Setting water applications:** Apply at a 1:100-1:200 dilution rate (128-64 fl. oz. per 100 gallons of transplant water or starter fertilizer). Make in-furrow or dibble application at the time of plant set. OxiDate 5.0 is chemically compatible with most water-soluble fertilizers, but conduct a compatibility test prior to mixing.

**Foliar applications immediately post planting:** Apply at a 1:100-1:200 dilution (128-64 fl. oz. for 100 gallons of water). Make application immediately following planting. Apply in a sufficient amount of water to achieve runoff to soil or plastic, typically 30 to 100 gallons of spray solution per acre.

**Foliar applications to established plants:** Follow: "Application Rates and Directions". When conditions favor development of disease, or at first sign/symptom of disease, apply at 1:100-1:200 (128-64 fl. oz. per 100 gallons of water) dilution. Maintain a 3-10 day spray schedule until control is achieved.

Always read and follow the product label for complete application instructions and restrictions. Conduct trial application on a representative area to confirm crop tolerance and compatibility.

\*Causal agent: Phomopsis obscurans
\*\*Causal agent: Neopestalotiopsis sp.



BioSafe Systems, LLC 22 Meadow Street East Hartford, CT 06108

www.BioSafeSystems.com 1-888-273-3088 Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.