



# OxiDate<sup>®</sup> 5.0

## FIFRA SECTION 2(ee) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

**ISSUE DATE:**

April 22, 2020

**EXPIRATION DATE:**

June 1, 2023

**FOR USE AND  
DISTRIBUTION ONLY  
IN THE STATE OF:**

AL, AR, CO, CT,  
DE, FL, GA, HI,  
IA, ID, IL, IN,  
KS, KY, LA, MA,  
MD, ME, MI, MN,  
MO, MS, MT, NC,  
ND, NE, NH, NJ,  
NM, NV, OH, OK,  
OR, PA, PR, RI,  
SC, SD, TN, UT,  
VA, VT, WA, WI,  
WV, WY



BioSafe Systems, LLC  
22 Meadow Street  
East Hartford, CT 06108  
www.biosafesystems.com  
1-888-273-3088

**EPA REGISTRATION NO.** 70299-28

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Crop: Corn:** field, sweet, popcorn

**Diseases:** Northern Corn Leaf Blight, Gray Leaf Spot, Bacterial Leaf Streak

**Application Rates and Directions****Curative application rates:**

1. For best results, apply at first sign of disease. Use OxiDate 5.0 at a dilution rate of 1:256 (50 fl. oz. per 100 gallons of clean water). Do not store and reuse mixed spray solution, prepare a fresh solution daily.
2. Spray plants in morning or evening. Apply in 10-500 gallons of water per acre, ensuring adequate coverage of upper and lower foliage, stems, branches and stalks. Final spray solution volume will depend on crop type, canopy size and/or growth stage.
3. Maintain a 3-10 day spray schedule until control is achieved.

**Rescue treatment rates:**

1. Concentrations up to 1:100 (1 gallon of OxiDate 5.0 for every 100 gallons of water) can be used as a rescue treatment for severe infestations.
2. Always test for phytotoxicity by spraying on few plants before using this rate on a large scale. It is recommended to use this rate outside of sensitive growth stages such as the bloom period.
3. Spray plants in morning or evening. Apply in 10-500 gallons of water per acre, ensuring adequate coverage of upper and lower foliage, stems, branches and stalks. Final spray solution volume will depend on crop type, canopy size and/or growth stage.
4. Maintain a 3-5 day spray schedule until control is achieved.

**Aerial Spray Treatments**

**Spray Drift Management** – Avoiding spray drift is the responsibility of the applicator.

Do not apply when wind conditions favor drift away from the intended area for treatment. Many factors including droplet size, equipment type and weather related factors determine the potential for spray drift.

To ensure optimum product performance, use at the foliar application rate indicated in sufficient water for adequate coverage of plant foliage. Apply between 3-20 gallons per acre of total spray solution. Do not make applications at a height greater than 10 ft. above the plant canopy, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to wind and evaporation. Do not exceed the maximum application rate or apply more often than labeled in the Application Instructions for that crop.

*See other diseases controlled by this product on the EPA product label.*

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.