



# OxiDate® 5.0

## FIFRA SECTION 2(ee) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

**ISSUE DATE:**

December 15, 2021

**EXPIRATION DATE:**

December 31, 2025

**FOR USE AND  
DISTRIBUTION ONLY  
IN THE STATE OF:**

**AL, AR, CO, CT,  
DE, FL, GA, HI,  
IA, ID, IL, IN,  
KS, KY, LA, MA,  
MD, ME, MI, MN,  
MO, MS, MT, NC,  
ND, NE, NH, NJ,  
NM, NV, OH, OK,  
OR, PA, PR, RI,  
SC, SD, TN, UT,  
VA, VT, WA, WI,  
WV, WY**

**EPA REGISTRATION NO.** 70299-28

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Crop: Cereal Grains & Commodities: Including, but not limited to:** Barley, Millet, Oats, Rice, Rye, Sorghum (Milo), Wheat, Wild Rice

**Diseases:** Bacterial Leaf Streak, Glume Blotch, Leaf Blight, Powdery Mildew, Bunt(s), Tan Spot

**Suppression of:** Fusarium Head Scab

**APPLICATION RATES AND DIRECTIONS**

**Curative:** 1:256 dilution. Apply at first sign/symptom of disease. Maintain a 3-10 day spray schedule until control is achieved.

**Rescue:** 1:100 dilution. Apply under severe disease conditions. Maintain a 3-5 day spray schedule until control is achieved. It is recommended to use this rate outside of sensitive growth stages such as the bloom period.

**AERIAL SPRAY TREATMENTS**

**Spray Drift Management** – Avoiding spray drift is the responsibility of the applicator.

Do not apply when wind conditions favor drift away from the intended area for treatment. Many factors including droplet size, equipment type and weather related factors determine the potential for spray drift.

To ensure optimum product performance, use at the foliar application rate indicated in sufficient water for adequate coverage of plant foliage. Apply between 3-20 gallons per acre of total spray solution. Do not make applications at a height greater than 10 ft. above the plant canopy, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to wind and evaporation. Do not exceed the maximum application rate or apply more often than labeled in the Application Instructions for that crop.

**See other crops and diseases controlled by this product on the EPA product label.**



BioSafe Systems, LLC

22 Meadow Street  
East Hartford, CT 06108

www.BioSafeSystems.com

1-888-273-3088

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.