



# OxiDate<sup>®</sup> 5.0

## FIFRA SECTION 2(ee) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

**ISSUE DATE:**

January 28, 2023

**EXPIRATION DATE:**

December 31, 2025

**FOR USE AND  
DISTRIBUTION ONLY  
IN THE STATE OF:  
CALIFORNIA**

**EPA REGISTRATION NO.** 70299-28

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Crop:** Celery

**Diseases:** Early and Late Blight

**Foliar Applications:** Follow **"Application Rates and Directions"**

**Curative:** 1:100 dilution. Apply at first sign/symptom of disease. Maintain a 3-10 day spray schedule until control is achieved.

**Application Rates and Directions**

OxiDate 5.0 works immediately on contact with any plant surface for control of plant diseases—see **Application Rates and Directions Chart**. Good coverage and wetting of the foliage is required. For increased coverage and penetration of spray, use a compatible non-ionic wetting agent/surfactant. For drift reduction and to aid spray deposition. Do not spray OxiDate 5.0 during conditions of intense heat, drought or poor plant vigor.

Run a plant sensitivity test when considering using higher spray concentrations greater than 0.39% v/v (1:256) by following instructions under "plant sensitivity testing". If plants show symptoms of phytotoxicity, decrease the spray solution concentration to a level that does not demonstrate symptoms.

1. For best results, apply at first sign of disease. Use OxiDate 5.0 at a dilution rate of 1:100 (1.0 gal. per 100 gallons of clean water). Do not store and reuse mixed spray solution, prepare a fresh solution daily.
2. Spray plants in morning or evening. Apply in 10-500 gallons of water per acre, ensuring adequate coverage of upper and lower foliage, stems, branches and stalks. Final spray solution volume will depend on crop type, canopy size and/or growth stage.
3. Maintain a 3-10 day spray schedule until control is achieved.



BioSafe Systems, LLC

22 Meadow Street  
East Hartford, CT 06108

www.BioSafeSystems.com

1-888-273-3088

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.