

## SPECIMEN LABEL

### FOR COMMERCIAL USE

EPA REGISTRATION NO. 70299-26

### ACTIVE INGREDIENTS:

Hydrogen Peroxide.....	10.00%
Peroxyacetic Acid.....	15.00%

OTHER INGREDIENTS:..... 75.00%

TOTAL:..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**STRONG OXIDIZING AGENT**  
**DANGER – PELIGRO**

*Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.*  
(If you do not understand this label, find someone to explain it to you in detail.)

### FIRST AID

#### If in eyes

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes.
- Call a poison control center or doctor for treatment advice.

#### If inhaled

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

#### If swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER: CORROSIVE.** Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Do not breathe vapor or spray mist. Wear a minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and a combination N, R or P filter; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved power air-purifying respirator with OV cartridges and combination HE filters. Wear goggles or face shield, rubber gloves, and protective clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or

using the toilet. Remove and wash contaminated clothing before reuse. Do not enter an enclosed area without proper respiratory protection.

### PHYSICAL AND CHEMICAL HAZARDS

**Corrosive.** Strong oxidizing agent. Do not use in concentrated form. Mix only with water in accordance with label instructions. Never bring concentrate in contact with other pesticides or oxidative agents.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Handlers who may be exposed to the undiluted product through mixing, loading, application, or other tasks must wear: coveralls over long-sleeved shirt and long pants, rubber gloves, chemical resistant footwear plus socks, protective eyewear (goggles, face shield or safety glasses). Handlers who may be exposed to the diluted product through application or other tasks must wear: long-sleeved shirt and long pants, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), notification to workers, and Restricted-Entry Interval (REI). The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

Handlers who may be exposed to the undiluted product through mixing, loading, application, or other tasks must wear: coveralls over long-sleeved shirt and long pants, rubber gloves, chemical resistant footwear plus socks, and protective eyewear goggles, face shield or safety glasses. Handlers who may be exposed to the dilute through application or other tasks must wear: long-sleeved shirt and long pants, and shoes plus socks.

For enclosed environments:

There is a Restricted Entry Interval (REI) of one (1) hour for this product when applied via spraying to growing plants, surfaces, equipment, structures and non-porous surfaces in enclosed environments such as glasshouses and greenhouses. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls worn over long-sleeved shirt and pants, waterproof gloves and shoes plus socks.

There is a Restricted Entry Interval (REI) of zero (0) hours for pre-plant dip, seed treatment, soil drench, mop, sponge, dip, soak, rinse or other non-spraying application methods when used in enclosed environments such as glasshouses and greenhouses.

For field applications:

Keep unprotected persons out of treated areas until sprays have dried. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls worn over long-sleeved shirt and pants, waterproof gloves and shoes plus socks.

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

**TREATMENT OF RAW, POST HARVEST FRUIT AND VEGETABLE AND PROCESSING WATERS**

**FOR REDUCTION AND CONTROL OF PATHOGENIC BACTERIA IN RAW, POST-HARVEST FRUIT AND VEGETABLE PROCESS AND WASH WATERS (REGULATED BY US EPA)**

Use SaniDate 15.0 to reduce (in 90 seconds) 99.9% of pathogenic bacteria *Escherichia coli* O157:H7, *Salmonella enterica*, and *Listeria monocytogenes* in processing waters used for washing fruits, and vegetables.

1. Add SaniDate 15.0 batch-wise or continuously to processing water without fruits, or vegetables present at a dilution of 0.8–1.9 fl. oz. per 25 gallons of water. This provides approximately 40–100 ppm of peroxyacetic acid and 27–64 ppm of hydrogen peroxide.
2. Allow the solution to circulate at least 90 seconds before adding raw fruits, or vegetables.
3. Adjust dose as needed to maintain a minimum product concentration of 40 ppm of peroxyacetic acid.
4. Allow for a minimum contact time of 90 seconds.
5. Prepare fresh process water daily. Do not reuse water that is badly fouled.
6. Contact your BioSafe Systems technical representative for specific applications.

**FOR TREATMENT OF RAW, POST-HARVEST FRUIT AND VEGETABLE PROCESSING WATERS TO CONTROL THE GROWTH OF NON-PUBLIC HEALTH MICROORGANISMS THAT CAN CAUSE SPOILAGE (REGULATED BY US EPA)**

1. Ensure that the water is recirculating or mixing in the processing tank or water line.

2. Prepare a use solution by diluting 1.5–1.9 fl. oz. of this product per 25 gallons of water. Ensure that the solution is thoroughly mixed. This provides 85–100 ppm of peroxyacetic acid and 57–67 ppm of hydrogen peroxide. Allow the solution to circulate at least 45 seconds before adding or treating raw fruits and vegetables.
3. Dose as needed to maintain 85–100 ppm of peroxyacetic acid by adding SaniDate 15.0 to processing water.
4. Allow a minimum contact time of 45 seconds.
5. Do not rinse.
6. Prepare a new solution daily to ensure effectiveness.
7. Contact your BioSafe Systems technical representative for specific applications.

**FOR TREATMENT OF RAW, UNPROCESSED FRUIT AND VEGETABLE SURFACES (REGULATED BY US EPA)**

Apply SaniDate 15.0 as a dip or spray to control the growth of non-public health microorganisms that may cause decay and/or spoilage on raw, post-harvest fruits and vegetables during the washing process. SaniDate 15.0 can be applied during physical cleaning processes, including at the roller spreader, washer manifold, and dip tank, on the brushes or elsewhere in the washing process prior to, simultaneously with or as a final rinse prior to packaging.

1. Prepare a use solution by diluting 1.5–1.9 fl. oz. of this product per 25 gallons of water. Ensure that the solution is thoroughly mixed. This provides 85–100 ppm of peroxyacetic acid and 57–67 ppm of hydrogen peroxide.
2. Apply the diluted sanitizing solution using a coarse spray directed at the fruits or vegetables, or by submerging the fruits or vegetables in the prepared solution.
3. Allow a minimum contact time of 45 seconds.
4. Do not rinse.
5. Contact your BioSafe Systems technical representative for specific applications.

**TREATMENT OF FRESH CUT AND PROCESSED FRUITS AND VEGETABLES AND PROCESSING WATERS**

**FOR TREATMENT OF FRESH CUT AND PROCESSED FRUIT AND VEGETABLE SURFACES TO CONTROL GROWTH OF NON-PUBLIC HEALTH MICROORGANISMS THAT CAN CAUSE SPOILAGE (REGULATED BY US FDA)**

Apply SaniDate 15.0 as a dip or spray to control the growth of non-public health microorganisms that may cause decay and/or spoilage on fresh cut and processed fruits and vegetables during the washing process.

1. Add 2.3 fl. oz. to 13.2 fl. oz. of this product to 100 gallons of water. This provides 30–175 ppm of peroxyacetic acid and 20–117 ppm of hydrogen peroxide. Ensure that the solution is thoroughly mixed.
2. Apply solution as a spray or dip. Allow a minimum contact time of 45 seconds.
3. Add this product to no more than 175 ppm residual peroxyacetic acid in the use solution in accordance with FCN #1554.
4. Contact your BioSafe Systems technical representative for produce specific rate recommendations.
5. A potable water rinse is not required after application of the diluted solution.

**SANITIZATION OF HARD, NON-POROUS FOOD CONTACT SURFACES**

For use in sanitizing of previously cleaned hard, non-porous food contact surfaces and equipment, processing and packaging equipment, boxing or packing equipment, such as pipelines, tanks, vats, fillers, evaporators, and pasteurizers. This product is not be used for sanitization of surfaces made of wood. Areas of use include:

- Meat And Poultry Slaughter and Further Processing Facilities
- USDA Inspected Food Manufacturing and Food Processing Plants
- Raw Fruit, Vegetable and Nut Packing Facilities
- Processed Fruit, Vegetable and Nut Processing Facilities
- Meat, Poultry, Seafood Processing/Packaging Plants

- Pet Food Manufacturing Plants
- Egg Packing and Egg Processing Plants
- Wineries, Breweries, Dairy, Beverage Bottling Plants
- Milk and Dairy Product Processing/Packing Plants
- Food Processing/Packing Plants
- Final Sanitizing Bottle Rinse
- Grocery, Supermarkets, Food Distribution and Storage Facilities

Before use in federally inspected meat and poultry food processing plants and dairies, food products must be removed from the room or covered to prevent contact with the sanitizing solution.

### Pathogenic Organisms

SaniDate 15.0 is an effective sanitizer against the following pathogenic organisms: *Escherichia coli*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Salmonella enterica*, *Escherichia coli* O157:H7, and *Campylobacter jejuni* on hard, non-porous food contact surfaces.

1. Pre-rinse surfaces and equipment to be sanitized with a warm potable water flush.
2. Wash surfaces and equipment with detergent or cleaning solution to remove gross particulate matter.
3. Rinse with potable water to remove suspended soils and residual detergent.
4. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
5. Fill closed systems with diluted sanitizer solution for a contact time of one (1) minute.
6. For open or not completely closed systems, apply sanitizing solution by using a coarse spray, mop/wipe or flood technique. Equipment may be immersed in the sanitizing solution.
7. Treated surfaces must remain visibly wet for at least one (1) minute.
8. Allow surfaces and equipment to drain thoroughly before resuming operations. Do not rinse.

### Non-Pathogenic Spoilage Organisms

SaniDate 15.0 is effective against non-pathogenic spoilage organisms, and yeasts.

1. Pre-rinse surfaces and equipment to be sanitized with a warm potable water flush.
2. Wash surfaces and equipment with detergent or cleaning solution to remove gross particulate matter.
3. Rinse with potable water to remove suspended soils and residual detergent.
4. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
5. Fill closed systems with diluted sanitizer solution for a contact time of one (1) minute.
6. For open or not completely closed systems, apply sanitizing solution by using a coarse spray, mop/wipe or flood technique. Equipment may be immersed in the sanitizing solution.
7. Treated surfaces must remain visibly wet for at least one (1) minute.
8. Allow surfaces and equipment to drain thoroughly before resuming operation. Do not rinse.

### SANITIZATION OF CONVEYORS, PEELERS, SLICERS, AND SAWS FOR MEAT, POULTRY, SEAFOOD, FRUITS, AND VEGETABLES

SaniDate 15.0 is an effective sanitizer against *Escherichia coli*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Salmonella enterica*, *Escherichia coli* O157:H7 and *Campylobacter jejuni*.

Use SaniDate 15.0 in the washing, rinsing and sanitizing of conveyor equipment, peelers, slicers, saws, corers, de-boners, dicers, knives, metal detectors, pasteurizers, peelers, scales, scrapers, collators, slicers, non-wooden cutting boards, tabletops, trays, pans, racks, platters, cans.

1. Remove edible products from equipment unless treating only the re-

turn portion of a conveyor.

2. Pre-rinse equipment and wash with detergent to lift away soils and gross particulate matter.
3. Rinse equipment with potable water to remove suspended soils and excess detergent.
4. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
5. Apply sanitizer solution to the return portion of the conveyor or to the equipment using a coarse spray or other means of wetting the surfaces. Control the volume of solution so as to permit maximum drainage and to prevent puddles. The conveyor surface may still be damp when food contact occurs.
6. Treated surfaces must remain visibly wet for at least one (1) minute.
7. Allow equipment to drain adequately before reusing; a dry surface is not required.

SaniDate 15.0 may be applied by flood applications, coarse spray, immersion, or circulation.

### FINAL SANITIZING BOTTLE RINSE

Use SaniDate 15.0 as a final sanitizing rinse for plastic, glass or metal returnable and non-returnable bottles/cans.

1. Wash bottles with detergent or cleaning solution and rinse with potable water.
2. Rinse bottles/cans with a solution prepared by mixing 0.41–0.94 fl. oz. of SaniDate 15.0 to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
3. Treated surfaces must remain visibly wet for at least one (1) minute.
4. Allow to drain dry.

### SANITIZING MILKING EQUIPMENT BY CLUSTER DIPPING

1. Clean the external surfaces of the milking systems after each use.
2. Manually, or automatically rinse and sanitize all system components using a SaniDate 15.0 sanitizing solution prepared by mixing 0.41–0.94 fl. oz. to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
3. Ensure solution fills clusters.
4. Treated surfaces must remain visibly wet for at least one (1) minute. Shake off well after dipping and allow to air dry. Do not rinse.

### FOR SANITIZING OF SHELL EGGS

1. To sanitize clean shell eggs intended for food or food products, spray with a use solution of 0.41–0.94 fl. oz. of SaniDate 15.0 to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide. The solution must be equal to or warmer than the eggs, but not to exceed 130° F.
2. Apply dilute solution as eggs are gathered as a coarse spray. Wet eggs thoroughly and allow to drain.
3. Treated surfaces must remain visibly wet for at least one (1) minute.
4. Eggs that have been sanitized with SaniDate 15.0 may be broken for use in the manufacture of egg products without a prior potable water rinse.
5. Eggs must be reasonably dry before casing or breaking.
6. The sanitizing solution must not be reused for sanitizing eggs.

### FOR SANITIZING OF HATCHING EGGS

1. Prepare a use solution by adding 0.41–0.94 fl. oz. SaniDate 15.0 to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
2. The solution must be equal to or warmer than the eggs, but not to exceed 130° F.
3. For hatching eggs apply the sanitizing solution as eggs are gathered or prior to setting using a coarse spray or flood so as to lightly wet egg shell surfaces.
4. Treated surfaces must remain visibly wet for at least one (1) minute.
5. Allow to drain dry.

## **SANITIZING HATCHERY ROOMS, POULTRY HOUSES, AND LIVESTOCK BUILDINGS**

1. Remove all animals and feed from premises, coops, crates and enclosures.
2. Remove gross soils, such as: litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Pre-rinse surfaces and equipment to be sanitized.
5. Thoroughly clean all surfaces with soap or detergent and rinse with water to remove detergent and suspended solids.
6. Prepare a use solution of SaniDate 15.0 by adding 0.41-0.94 fl. oz. to 5 gallons of potable water. This provides 109-250 ppm peroxyacetic acid and 73-167 ppm of hydrogen peroxide.
7. Apply the SaniDate 15.0 solution to surfaces using a coarse spray, mop/wipe or flood.
8. Immerse all types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure in the SaniDate 15.0 solution.
9. Treated surfaces must remain visibly wet for at least one (1) minute.
10. Allow surfaces and equipment to drain adequately before use. Do not rinse.
11. Ventilate buildings, coops and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed, set or dried.

## **PACKINGHOUSE SANITIZATION**

SaniDate 15.0 is an effective sanitizer against the following microorganisms: *Xanthomonas campestris (axonopodis) pv. citrumelo* (citrus canker surrogate), as well as *Escherichia coli*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Salmonella enterica*, *Escherichia coli* O157:H7 and *Campylobacter jejuni*.

1. Pre-rinse surfaces and equipment to be sanitized.
2. Remove gross contamination with a cleaner or other suitable detergent and rinse with potable water.
3. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
4. Use as a general sanitizing coarse spray to reduce bacterial contamination of walls, floors, conveyers and harvesting containers.
5. Equipment may be immersed in the SaniDate 15.0 solution.
6. Treated surfaces must remain visibly wet for at least one (1) minute.
7. Allow to air dry, do not rinse.

## **HARVESTING AND FIELD EQUIPMENT AND TRANSPORTATION VEHICLE SANITIZATION**

Use SaniDate 15.0 to sanitize harvest equipment such as pickers, trailers, trucks (including truck body parts and tires), bins, packing crates, ladders, power tools, hand tools, gloves, rubber boots, pruning shears or other equipment that may transfer *Xanthomonas campestris (axonopodis) pv. citrumelo* (citrus canker surrogate). This product can also be used to sanitize surfaces contaminated with *Escherichia coli*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Salmonella enterica*, *Escherichia coli* O157:H7 and *Campylobacter jejuni*.

1. Remove gross contamination with a cleaner or other suitable detergent and rinse with water.
2. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
3. Apply as a general sanitizing coarse spray.
4. Treated surfaces must remain visibly wet for at least one (1) minute.
5. Allow to air dry, do not rinse.

## **SURFACES TREATED TO CONTROL THE SPREAD OF CITRUS CANCKER**

Use SaniDate 15.0 to control the spread of citrus cancker between inanimate

surfaces and inanimate surfaces to plants. This product is for sanitizing surfaces such as packinghouse conveyers and harvesting equipment and containers. This product is not for treatment of infected plants.

## **SANITIZATION OF HARD, NON-POROUS, NON-FOOD CONTACT SURFACES**

SaniDate 15.0 is an effective hard, non-porous, non-food contact surface sanitizer against *Staphylococcus aureus*, and *Klebsiella pneumoniae*. SaniDate 15.0 may be applied by flood applications, coarse spray, immersion, or circulation. Sanitize surfaces including ceilings, drain grates, floors, and walls. This product is not be used for sanitization of surfaces made of wood.

1. Pre-rinse surfaces and equipment to be sanitized with potable water.
2. Remove gross contamination with a cleaner or other suitable detergent and rinse with potable water.
3. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
4. Apply by wiping, mopping, coarse spray, or flood. Equipment may be immersed in sanitizing solution.
5. Treated surfaces must remain visibly wet for five (5) minutes.
6. Allow items and/or surfaces to air dry before resuming operation.
7. Do not rinse.

## **SANITIZATION OF HARD, NON-POROUS, NON-FOOD CONTACT PACKAGING EQUIPMENT**

1. Pre-rinse equipment to be sanitized with potable water.
2. Wash equipment with detergent or cleaning solution to remove gross particulate matter.
3. Rinse equipment with potable water to remove suspended soils and excess detergent.
4. Prepare a use solution by adding 0.41–0.94 fl. oz. of this product to 5 gallons of potable water. This provides 109–250 ppm peroxyacetic acid and 73–167 ppm of hydrogen peroxide.
5. Apply by spray, or circulation techniques as appropriate. Allow a contact time of at least five (5) minutes.
6. For open or not completely closed systems, Apply SaniDate 15.0 by immersion, coarse spray, mop/wipe or flood technique. Treated surfaces must remain visibly wet for at least five (5) minutes.
7. Allow equipment and/or surfaces to air dry before resuming operation.
8. Do not rinse.

## **GENERAL DISINFECTION**

SaniDate 15.0 is an effective one-step cleaner and disinfectant against gram positive and negative bacteria (vegetative forms): *Staphylococcus aureus*, *Salmonella enterica*, *Pseudomonas aeruginosa*. It is effective in hard water (up to 400 ppm as calcium carbonate equivalent), on lightly soiled surfaces. Use SaniDate 15.0 in general commercial and medical environments to clean, disinfect, and deodorize hard, non-porous surfaces. This product is not be used for the disinfection of surfaces made of wood.

### **Use SaniDate 15.0 on the following surfaces:**

Floors, walls, drains and other non-porous surfaces such as tables, chairs, counter tops, garbage cans/bins, bathroom fixtures, sinks, bed frames, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, sealed asphalt, glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel or glass.

### **Areas of use include:**

- Hospitals, surgical and obstetrical suites; operating tables, housekeeping services; physical therapy departments; nursing homes, health care facilities, autopsy facilities, pharmaceutical and chemical processing facilities and equipment.
- Schools, colleges, industrial facilities, dietary areas, office buildings, recreational facilities, retail and wholesale establishments.
- Animal hospitals, veterinary clinics, animal life science laboratories, kennels, kennel runs, cages, feeding and watering equipment, pet shops, zoos, pet animal quarters, poultry premises, trucks, hatcheries,

and livestock quarters.

- Dairies, wineries, breweries, and beverage plants
- Packinghouses, food processing, food distribution and storage, beverage processing facilities, milking parlors, dairy production and transfer facilities and equipment
- Farms, farm equipment and harvesting equipment
- USDA inspected food manufacturing and food processing plants
- Meat and meat products processing, packing, and rendering plants
- Meat and poultry slaughter and further processing plants
- Milk and dairy processing/packing plants
- Egg processing/packing plants
- Seafood and poultry processing/packing plants
- Raw/Processed fruit, vegetable and nut processing/packing plants
- Grocery stores, supermarkets, food distribution and storage facilities
- Eating establishments

#### **Disinfection:**

1. For moderately soiled surfaces a pre-cleaning step is not required. For grossly contaminated surfaces, pre-rinse surfaces and equipment to be disinfected with potable water.
2. Wash surfaces and equipment with detergent or cleaning solution to remove gross filth. Rinse with potable water to remove suspended soils and residual detergent.
3. Prepare a disinfecting solution by mixing 1.5–8.7 fl. oz. of SaniDate 15.0 in 5 gallons of water. This provides 393–2,300 ppm of peroxyacetic acid and 262–1,522 ppm of hydrogen peroxide.
4. Apply the SaniDate 15.0 solution with a mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking to wet all surfaces thoroughly. Treated surfaces must remain visibly wet for a minimum of ten (10) minutes.
5. Rinse all treated surfaces that will contact food or commodities with potable water before use.

#### **DISINFECTING HOSPITALS, DENTAL OFFICES, NURSING HOMES, AND OTHER HEALTH CARE INSTITUTIONS**

For disinfecting hard, non-porous surfaces such as floors, walls, countertops, bathing areas, lavatories, bed frames, tables, chairs, and garbage pails. For heavily soiled surfaces, a pre-cleaning step is required.

1. Remove gross contamination with a cleaner or other suitable detergent and rinse with water.
2. Add 1.5–8.7 fl. oz. of SaniDate 15.0 to 5 gallons of water. This provides 393–2,300 ppm of peroxyacetic acid and 262–1,522 ppm of hydrogen peroxide.
3. Apply the SaniDate 15.0 solution with a mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking to wet all surfaces thoroughly.
4. Treated surfaces must remain visibly wet for a minimum of ten (10) minutes.

#### **DISINFECTING PHARMACEUTICAL AND COSMETIC SURFACES**

SaniDate 15.0 is recommended for use on hard, non-porous, environmental surfaces such as floors, walls and processing equipment in pharmaceutical and cosmetic processing facilities.

1. Remove gross contamination with a cleaner or other suitable detergent and rinse with water.
2. Add 1.5–8.7 fl. oz. of SaniDate 15.0 to 5 gallons of water. This provides 393–2,300 ppm of peroxyacetic acid and 262–1,522 ppm of hydrogen peroxide.
3. Apply the SaniDate 15.0 solution with a mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking to wet all surfaces thoroughly.
4. Treated surfaces must remain visibly wet for a minimum of ten (10) minutes.

#### **DISINFECTING HAIRDRESSING SALON/BARBER SHOP INSTRUMENTS AND TOOLS**

Immerse pre-cleaned barber/salon tools (combs, brushes, razors, manicure/

pedicure tools, clippers, scissors, trimmer blades) in a solution of SaniDate 15.0 at a dilution of 0.3 fl. oz. per gallon of water, for at least ten (10) minutes. This provides 393 ppm of peroxyacetic acid and 262 ppm of hydrogen peroxide. Rinse instruments thoroughly and dry before reuse. A fresh use-solution should be prepared daily or more often if the use solution becomes cloudy or soiled. Note: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried and kept in a clean non-contaminated receptacle. Prolonged immersion may cause damage to stainless steel or metal instruments.

#### **FIELD AND FARM EQUIPMENT DISINFECTION**

SaniDate 15.0 may be used to disinfect hard, non-porous harvest equipment such as pickers, harvesters, tractors, trailers, forklifts, trucks (including truck body parts and tires), bins, packing crates, ladders, power tools, hand tools, gloves, rubber boots, pruning shears or other equipment.

1. Remove gross contamination with a cleaner or other suitable detergent and rinse with water.
2. Apply SaniDate 15.0 at a rate of 0.3 fl. oz. per gallon of water as a general coarse spray. This provides 393 ppm of peroxyacetic acid and 262 ppm of hydrogen peroxide.
3. Treated surfaces must remain visibly wet for ten (10) minutes.
4. Allow to air dry, do not rinse.

#### **TRACTOR TRAILER AND TRANSPORTATION VEHICLES DISINFECTION**

SaniDate 15.0 may be used to disinfect and deodorize hard, non-porous equipment such as trucks, trailers, cabs, crates, (including truck body parts and tires, mats, wheels). Use SaniDate 15.0 to prevent cross contamination of bacteria across treated surfaces between loads.

1. Remove gross contamination with high pressure water and cleaner or other suitable detergent and rinse with water.
2. Apply SaniDate 15.0 at a rate of 0.3 fl. oz. per gallon of water as a general coarse spray. This provides 393 ppm of peroxyacetic acid and 262 ppm of hydrogen peroxide.
3. Treated surfaces must remain visibly wet for ten (10) minutes.
4. Rinse all treated surfaces that will contact food or commodities with potable water before use.

#### **ANIMAL HEALTH**

SaniDate 15.0 can be used in animal hospitals, animal laboratories, kennels, pet shops, zoos, pet animal quarters, poultry premises, poultry hatcheries, livestock and dairy quarters. When used as directed, it is specifically designed to disinfect, deodorize and clean inanimate, hard, non-porous surfaces such as walls, floors, sink tops, furniture, operation tables, kennel runs, cages and feeding and watering equipment. In addition, it will disinfect bins and cans, and any other hard, non-porous areas that are prone to odors caused by microorganisms.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

#### **DISINFECTION OF POULTRY HOUSES, ANIMAL HOUSING, LIVESTOCK BUILDINGS AND HATCHERIES**

1. Remove all poultry, livestock and feed from premises, coops, crates and enclosures.
2. Remove gross soils, such as litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Pre-rinse surfaces and equipment to be disinfected.
5. Thoroughly clean all surfaces with soap or detergent and rinse with water.
6. Apply SaniDate 15.0 at a rate of 1.5–8.7 fl. oz. of SaniDate 15.0 in 5 gallons of water to surfaces and equipment using a course spray or flood. This provides 393–2300 ppm of peroxyacetic acid and 262–1,522 ppm of hydrogen peroxide.

7. Immerse all types of equipment used in handling and restraining poultry, as well as forks, shovels and scrapers used for removing litter and manure.
8. Treated surfaces must remain visibly wet for a minimum of ten (10) minutes.
9. Ventilate buildings, coops and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed, set or dried.
10. Thoroughly rinse all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

#### **TERRARIUM AND SMALL ANIMAL CAGE DISINFECTION**

1. Remove all animals and feed from enclosure to be cleaned.
2. Thoroughly clean all hard, non-porous surfaces with soap or detergent and rinse with water.
3. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with a solution of 0.3 fl. oz. of this product per gallon of water. This provides 393 ppm of peroxyacetic acid and 262 ppm of hydrogen peroxide. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet.
4. Treated surfaces must remain visibly wet for a minimum of ten (10) minutes.
5. Thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.
6. Do not return animals to the habitat until it is dry and ventilated.
7. Clean terrarium at least once weekly or more as needed.

#### **FOOT BATH MATS, PADS, WALK THROUGH TRAYS**

Place foot bath mats, pads or trays at the entrances of all rooms and buildings to prevent cross contamination across treated surfaces in animal containment areas, livestock and dairy quarters, poultry premises, greenhouses, packing houses, food processing and rendering plants.

1. Prior to use of this product, rinse or brush footwear surfaces to remove gross filth.
2. Make a solution using 0.3–0.4 fl. oz. of SaniDate 15.0 per gallon of water (393–525 ppm of peroxyacetic acid) and add to foot bath mat, pad or tray, filling to capacity.
3. Place boots and shoes in the foot bath mat, pad or tray containing the recommended solution of SaniDate 15.0. Allow treated surfaces to remain visibly wet for ten (10) minutes prior to entering next area. Change solution daily or as needed.

#### **DISINFECTION OF HARD, NON-POROUS FOOD-CONTACT SURFACES IN FOOD PROCESSING PLANTS AND FOOD SERVICE ESTABLISHMENTS**

Before using this product, food products and packaging materials must be removed from area or carefully protected.

1. Prior to use of this product, remove gross soil particles from surfaces to be treated. For heavily soiled surfaces, a pre-wash is required.
2. Apply 0.3 fl. oz. of SaniDate 15.0 per gallon of water with a mop, cloth, sponge, or hand trigger spray to wet all surfaces thoroughly. This provides 393 ppm of peroxyacetic acid and 262 ppm of hydrogen peroxide.
3. Treated surfaces must remain visibly wet for ten (10) minutes.
4. Rinse all treated surfaces thoroughly with potable water before operations are resumed.

#### **PACKINGHOUSE, FOOD STORAGE FACILITIES, FOOD PROCESSING AND RENDERING PLANT DISINFECTION**

Apply SaniDate 15.0 on all hard, non-porous surfaces and equipment found in commercial packinghouses including dump tanks, drenches, crates, containers, conveyors, storages, walls, floors, and process lines.

#### **For Pre-Cleaned Surfaces:**

Use SaniDate 15.0 for hard, non-porous surfaces that are lightly soiled or have been pre-cleaned to remove gross contamination.

1. Cover or remove all food and packaging materials before disinfection.
2. Apply 0.3 fl. oz. of SaniDate 15.0 per gallon of water with a mop, cloth, sponge, or hand trigger spray to wet all surfaces thoroughly. This provides 393 ppm of peroxyacetic acid and 262 ppm of hydrogen peroxide.
3. Treated surfaces must remain visibly wet for ten (10) minutes.
4. Rinse all treated surfaces thoroughly with potable water before operations are resumed.

#### **CLEANING POULTRY, SWINE, LIVESTOCK WATER SYSTEMS WHEN THE SYSTEM IS NOT IN USE**

To remove scale, calcium, iron, magnesium, heavy soils, polysaccharides and deposits from vitamins and medications from livestock watering systems use SaniDate 15.0 at 0.15–0.30 fl. oz. per gallon of water. This provides 197–393 ppm of peroxyacetic acid and 131–262 ppm of hydrogen peroxide. When used as directed, SaniDate 15.0 will remove organic and inorganic deposits that reduce water flow and clog nipples. Allow system to run for 6–24 hours depending on the conditions. Following the cleaning process, rinse with potable water to remove the cleaning solution from the watering line, nipples and cups. Never mix SaniDate 15.0 with any other product.

#### **CLEANING POULTRY, SWINE, LIVESTOCK WATERING OPERATING SYSTEMS (WHEN ANIMALS ARE PRESENT)**

After water lines have been cleaned, use SaniDate 15.0 at 0.3–0.45 fl. oz. per 100 gallons of water to control mineral build up in watering lines. This provides 3.9–5.8 ppm of peroxyacetic acid and 2.6–3.9 ppm of hydrogen peroxide. Never use SaniDate 15.0 more than 5 consecutive days to clean the operating system. Never mix SaniDate 15.0 with any other product. If cleaning the operating system, stop the use of SaniDate 15.0 twenty-four (24) hours prior to vaccination or medication via the water line.

#### **CONTROL OF NON-PUBLIC HEALTH SLIME-FORMING BACTERIA AND BIOFOULING IN COOLING WATER SYSTEMS (COOLING TOWERS, EVAPORATIVE CONDENSERS, AIR WASHERS) AND ORNAMENTAL WATER FEATURES**

1. Severely fouled systems should be cleaned before adding the SaniDate 15.0 solution. SaniDate 15.0 should be added in the water system directly, and not mixed with any other chemicals or additives. Never add SaniDate 15.0 into any feeding device, such as shot feeders, filter housings, by-pass feeders, or miscellaneous piping of any kind, because dangerous acute decomposition can occur. Discontinue the use of chlorine or bromine products prior to using SaniDate 15.0. Contamination with other chemicals could result in product decomposition.
2. Add SaniDate 15.0 only to water at a point in the system where uniform mixing and even distribution will occur.
3. For shock (slug) treatment for moderately to severely fouled systems: Add 5–20 fl. oz. of this product per 1,000 gallons of process water (7–27 ppm peroxyacetic acid, 5–18 ppm hydrogen peroxide). Repeat as necessary until microbiological control is evident. Thereafter, to maintain control use 1.5–7.5 fl. oz. of SaniDate 15.0 per 1,000 gallons of process water (2–10 ppm of peroxyacetic acid) as a continuous treatment method.
4. Continuous dosing methods usually require 1.5–5 fl. oz. per 1,000 gallons of water (2–7 ppm peroxyacetic acid, 1–5 ppm hydrogen peroxide) to achieve adequate results.
5. Intermittent dosing treatments usually require dose cycles of a minimum once per day, up to 6 times per 24 hours. Recommended rates for intermittent dose cycles are 5–10 fl. oz. of SaniDate 15.0 per 1,000 gallons of process water (7–14 ppm peroxyacetic acid, 5–9 ppm hydrogen peroxide).

**Air Washers:** SaniDate 15.0 may be used to control bacteria and biofouling in industrial air washing/scrubbing systems. The air washer must have operational and effective mist elimination systems. Prior to use of this product, heavily fouled systems must be pre-cleaned using an appropriate cleaner. Continuous dosing methods will require 2–7 ppm and intermittent dosing methods require 7–14 ppm (as peroxyacetic acid) depending on the type of system and the level of microbiological control desired.

## TREATMENT OF GREENHOUSE IRRIGATION WATER TO CONTROL SLIME-FORMING BACTERIA

For treatment of irrigation water, use a dilution rate of 1:51,364–1:6,519 (2.50–19.6 fl. oz. of SaniDate 15.0 per 1,000 gallons of water; equivalent to 3–26 ppm of peroxyacetic acid).

## TREATMENT OF AGRICULTURAL IRRIGATION WATER AND DRAINAGE DITCHES

Use SaniDate 15.0 at the following rates to suppress/control bacteria in irrigation water and drainage ditches.

- Bacteria: 3–100 fl. oz. per 1,000 gallons of water (1:50,960–1:1,274 dilution)

Product can be simply added to the body of water.

## TREATMENT OF AGRICULTURAL WATER USED FOR PESTICIDE SPRAY SOLUTIONS

Use SaniDate 15.0 to treat and suppress bacteria in water collected from open or closed sources including but not limited to wells, streams, ponds, lakes, canals, irrigation reservoirs, and drainage ditches, used for pesticide spray solutions and mixtures. Add SaniDate 15.0 at the recommended rate to water in spray or mix tank. Mix and allow a contact time of 3–5 minutes ***before adding other pesticides to spray solution.***

- Bacteria: 3–100 fl. oz. per 1,000 gallons of water (1:50,960–1:1,274 dilution)

### Tank Mixing Instructions

**1. Before adding other pesticides to the spray solution:** Mix the SaniDate 15.0 solution first and allow a contact time of 3–5 minutes.

**2. When used with Conventional Bactericides/Fungicides/Insecticides/Miticides:** It is recommended to not exceed a rate of 0.026% v/v (1:3,822; 0.033 fl. oz. per gallon or 3.3 fl. oz. per 100 gallons of water) with SaniDate 15.0. This recommendation applies to pesticides with or without metal ion(s).

**3. When used with Organic (Biorational/Botanical/Biological) Bactericides/ Fungicides/ Insecticides:** It is recommended to not exceed a rate of 0.013% v/v (1:7,644; 0.017 fl. oz. per gallon or 1.7 fl. oz. per 100 gallons of water) with SaniDate 15.0 Spray Tank Water treatment can be used with Biorational/Botanical based Bactericides/ Fungicides/Insecticides/Miticides (Ex. Neem Oil, Sulfur, Plant Extracts etc.), *Bacillus* based Bio-Fungicides (spore containing or spent fermented media), Bt based Bio-Insecticides and Copper based Bactericides/Fungicides ***Do not use SaniDate 15.0 with Mycoinsecticides (*Beauveria*, *Metarhizium*, *Isaria* based) or with other biological active ingredients not listed above.***

**4. When used with Micro-Foliar Fertilizers:** It is recommended to not exceed a rate of 0.026% v/v (1:3,822; 0.033 fl. oz. per gallon or 3.3 fl. oz. per 100 gallons of water) with SaniDate 15.0.

### Compatibility:

SaniDate 15.0 is compatible as a direct injection or tank-mix with many commonly used pesticides, fertilizers, adjuvants and non-ionic surfactants but has not been fully evaluated with all of these. Do not direct inject or tank mix SaniDate 15.0 into the irrigation system or in spray tank with pesticides, surfactants or fertilizers without conducting a compatibility test to show it is physically compatible, effective and non-injurious under your use conditions. Compatibility test is also recommended when tank mixing SaniDate 15.0 with copper or other pesticides containing metals at a dilution rate stronger than 1:3822.

To ensure compatibility, evaluate them prior to use as follows: Using a suitable container, add proportional amounts of product to water. Add wettable powders first, followed by water dispersible granules, then by liquid flowables and lastly, emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible. Test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Contact your BioSafe

Systems representative for further instructions.

## TREATMENT OF WATER DRAWN FROM OPEN AND CLOSED WATER SOURCES USED FOR DUST ABATEMENT

Use SaniDate 15.0 at the following rates to suppress/control bacteria in water drawn from open and closed water sources such as wells, streams, ponds, reservoirs, irrigation canals, irrigation water and drainage ditches used to control dust on unpaved gravel and dirt roads.

- Bacteria: 3–100 fl. oz. per 1,000 gallons of water (1:50,960–1:1,274 dilution)

Prepare the mixture at least 3–5 minutes prior to application for dust abatement. Apply to the road surface using a water truck (or tractor or spraying device) with equipped with a watering system.

## TREATMENT OF AGRICULTURAL IRRIGATION WATER USED FOR FRUIT, VEGETABLE AND ROW CROPS

Use SaniDate 15.0 to treat irrigation water during all phases of crop production including pre-plant irrigation and throughout the crop cycle to suppress/control bacteria in irrigation water used for fruit, vegetable and row crop production. SaniDate 15.0 can be used up to and including the day of harvest.

- Bacteria: 3–100 fl. oz. per 1,000 gallons of water (1:50,960–1:1,274 dilution)

Apply this product as a direct injection into the water at the point of intake and applied through a sprinkler system (including solid set, center pivot, lateral move, end tow, side wheel roll, traveling big gun or hand move), drip/micro irrigation system, flood (basin), or furrow. For best results, treat water every time crop is irrigated or at a minimum during the last 2–3 irrigations prior to harvest.

## FOR TREATMENT OF IRRIGATION WATER TO SUPPRESS/CONTROL BACTERIA

For treatment of irrigation water, use a dilution rate of 1:50,960–1:1274 (3–100 fl. oz. of SaniDate 15.0 per 1,000 gallons of water; equivalent to 4–134 ppm of peroxyacetic acid).

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original containers in a cool, well-ventilated area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If wastes cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

#### For non-refillable containers equal to or less than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

**For non-refillable containers greater than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the

container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

**For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**WARRANTY  
CONDITIONS OF SALE AND LIMITATION  
OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of BIOSAFE SYSTEMS, LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold BIOSAFE SYSTEMS, LLC and Seller harmless for any claims relating to such factors to the extent consistent with applicable law.

BIOSAFE SYSTEMS, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above when used in accordance with directions under normal use conditions. To the extent consistent with applicable law. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or BIOSAFE SYSTEMS, LLC and Buyer and User assume the risk of any such use. BIOSAFE SYSTEMS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall BIOSAFE SYSTEMS, LLC or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF BIOSAFE SYSTEMS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF BIOSAFE SYSTEMS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

BIOSAFE SYSTEMS, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of BIOSAFE SYSTEMS, LLC.



For additional information on SaniDate® 15.0, call us toll-free at 1.888.273.3088 or visit [www.biosafesystems.com](http://www.biosafesystems.com).

©2020 BioSafe Systems, LLC. SaniDate 15.0® is a registered trademark of BioSafe Systems, LLC. Always read and follow label directions.

**V11-030420** 12.20